


SACROILIAC JOINT INJECTION




For Low Back and Buttocks Pain

A sacroiliac joint injection is an outpatient procedure diagnosing and treating low back and buttocks pain.

What is the sacroiliac joint and why is a sacroiliac joint injection helpful?

The sacroiliac joint is a large joint in your lower back and buttocks region. When the joint becomes painful, it can cause pain in its immediate region or it can refer pain into your groin, abdomen, hip, buttock or leg. (see image at left.)

A sacroiliac joint injection serves several purposes. First, by placing numbing medicine into the joint, the amount of immediate pain relief you experience will help confirm or deny the joint as a source of your pain. That is, if you obtain complete relief of your main pain while the joint is numb it means this joint is more likely than not your pain source. Furthermore, time-release cortisone will be injected into the joint to reduce any presumed inflammation, which on many occasions can provide long-term pain relief.





What will happen to me during the procedure?

If needed, an IV will be started so that adequate relaxation medication can be given. After lying on an x-ray table, the skin over your lower back/buttock will be well cleansed. Next, the physician will numb a small area of skin with medicine (anesthetic) which stings for a few seconds. The physician then will use x-ray guidance to direct a very small needle into the joint. He will then inject several drops of contrast dye to confirm that the medication only goes into the joint. A small mixture of numbing medication (anesthetic) and anti-inflammatory cortisone will then be slowly injected.

What should I do and expect after the procedure?

20-30 minutes after the procedure, you will move your back to try to provoke your usual pain. You will report your remaining pain, (if any) and also record the relief you experience during the next week, on a "pain diary" we will provide. You may or may not obtain improvement in the first few hours after the injection, depending on if the sacroiliac joint is your main pain source. The pain diary is an important component of your care. It helps your treating physician to be informed of your results so future tests and/or needed treatment can be planned.





How will I feel after the procedure?

On occasion, the part of your treated spine may feel slightly weak or have an odd feeling in your leg for a few hours after the injection. You may notice a slight increase in your pain lasting for several days, as the numbing medicine wears off before the cortisone becomes effective.

Ice is typically more helpful than heat during the first 2-3 days after the injection.

You may begin to notice an improvement in your pain 2-5 days after the injection. If you do not notice improvement within 10 days after the injection, it is unlikely to occur.


You may take your regular medications after the procedure, but try to limit any pain medications for the first 4-6 hours after the procedure. This will ensure that the diagnostic information obtained from the procedure is accurate.

You may be referred for physical, chiropractic treatment, or massage therapy after the injection while the numbing medicine is effective and/or over the several weeks while the cortisone is working.

When can I resume activity?

On the day of the injection you should not drive and should avoid any strenuous activities. On the day after the procedure, you may return to your regular activities.

Sacroiliac Joint Injections



UTAH PAIN RELIEF INSTITUTE
Your ~~Pain~~ Solution Center